

Technical University of Denmark



Quantifying Urban Foodprints and Mitigation Opportunities

Goldstein, Benjamin Paul; Fernandez, John; Birkved, Morten; Hauschild, Michael Zwicky

Publication date:
2015

[Link back to DTU Orbit](#)

Citation (APA):

Goldstein, B. P., Fernandez, J., Birkved, M., & Hauschild, M. Z. (2015). Quantifying Urban Foodprints and Mitigation Opportunities. Poster session presented at 8th Conference of the International Society for Industrial Ecology, Guildford, United Kingdom.

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Quantifying Urban Foodprints and Mitigation Opportunities

Benjamin Goldstein^{1,2} (bgol@dtu.dk and ) , John Fernandez², Morten Birkved¹ and Michael Hauschild¹

¹Technical University of Denmark, Quantitative Sustainability Assessment Division

²Massachusetts Institute of Technology, Department of Architecture - Urban Metabolism Group

Introduction

As centers of wealth and population, cities are and will continue to be the major driving force of anthropogenic environmental impacts. Overviews of urban transport [1], building energy [2] and water consumption [3] have shown the importance of these activities in influencing urban environmental integrity. Urban food consumption has not received a similar treatment despite its large contributions to global climate change [4], non-renewable resource consumption [5] and other ecological challenges. This study attempts to fill that gap by asking:

What is food consumption's role in a city's environmental profile (the 'foodprint') and how can urban designers re-imagine cities to mitigate it?

Methods

Comprehensive review of urban metabolism and related literature to find quantified urban foodprints in terms of:

- Urban food demand (tons/cap/a) - *MFA*
- Ecological footprint (global ha/cap/a) - *EF*
- Carbon footprint (tons CO₂ eq./cap/a) - *CF*




Fig 1: Overview of research methods




Fig 2: Map showing the locations of foodprint studies included in the review

Acknowledgements

Thanks to Otto Mønsted Foundation (grant: 14-70-1109), Augustinus Fund (grant: 14-3738) and Reinholdt W. Jorck og Hustrus Fund (grant: 14-DI-0434) for their generous financial support of this research.

Results

The urban foodprint is a significant driver of the reviewed cities' overall environmental pressures at (figure 3):

- 1 ton food/cap/a (figure 4a)
- 1.2 global hectare/cap/a (figure 4b)
- 1.9 tons CO₂ eq./cap/a (figure 4c)

Plotting against per capita GDP shows moderate positive correlation with foodprint and food waste hinting at consumption patterns that will increase in terms of embodied impacts (more animal products intake) and wastefulness with future urbanization [6] (figure 4d).




Fig 3: Rank of foodprint relative to other major urban metabolic drivers (e.g. transport, building energy, etc.)




Fig 4a: Per-capita mass foodprint vs. per-capita GDP




Fig 4b: Per-capita carbon foodprint vs. per-capita GDP




Fig 4c: Per-capita ecological foodprint vs. per-capita GDP




Fig 4d: Per-capita waste foodprint vs. per-capita GDP

Conclusions

The urban foodprint is significant in scale and is poised to grow lockstep with future urbanization. Urban food waste is also set to grow simultaneously, which combined with the current linear metabolism of many cities food systems could result in a significant nutrient exhaust and methane production. These will challenge urban designers and policy makers to explore new infrastructural forms and policies for future cities:

- *Circular metabolic strategies*: wastewater sludge recycling, composting;
- *Behavioral nudging*: food waste attenuation, lower and dairy intake;
- *Alternative food supply-chains*: urban and peri-urban agriculture.

The combined effect of such interventions on the urban foodprint is unknown, making it a ripe question of practical importance for future urban metabolism research.

Works Cited

- [1] Grubler, A., X. Bai, T. Buettner, S. Dhakal, D.J. Fisk, T. Ichinose, J.E. Keirstead, et al. 2012. Chapter 18 - Urban Energy Systems. In *Global Energy Assessment - Toward a Sustainable Future*, 1307–1400. Cambridge University Press, Cambridge, UK and New York, NY, USA and the International Institute for Applied Systems Analysis, Laxenburg, Austria.
- [2] Darrel Jenerette, G. and L. Larsen. 2006. A global perspective on changing sustainable urban water supplies. *Global and Planetary Change* 50(3-4): 202–211.
- [3] Kenworthy, J. and F. Laube. 1996. Automobile Dependence in Cities: An International Comparison of Urban Transport and Land Use Patterns with Implications for Sustainability. *Environmental Impact Assessment Review* 16(4-6): 279–308.
- [4] Foley, J. a, N. Ramankutty, K. a Brauman, E.S. Cassidy, J.S. Gerber, M. Johnston, N.D. Mueller, et al. 2011. Solutions for a cultivated planet. *Nature* 478(7369): 337–42.
- [5] FAO. 2006. *Livestock's Long Shadow*. Rome, IT. <http://www.fao.org/docrep/010/a0701e/a0701e00.HTM>.
- [6] Goldstein, B., M. Birkved, J. Fernandez and M. Hauschild. 2015. Surveying the environmental footprint of urban food consumption (forthcoming)